arXiv:hep-th/9608077v1 12 Aug 1996

UFIFT-HEP-96-

August 12, 1996

Superalgebras in $N = 1$ Gauge Theories

Pierre Ramond

Department of Physics

University of Florida

Gainesville FL 32611

ABSTRACT:

$N = 1$ supersymmetric gauge theories with global flavor symmetries contain a gauge invariant W-superalgebra which acts on its moduli space of gauge invariants. With adjoint matter, this superalgebra reduces to a graded Lie algebra. When the gauge group is $SO(n_c)$, with vector matter, it is a W-algebra, and the primary invariants form one of its representation. The same superalgebra exists in the dual theory, but its construction in terms of the dual fields suggests that duality may be understood in terms of a charge conjugation within the algebra. We extend the analysis to the gauge group E_6.

Consider a $N=1$ supersymmetric gauge theory with a simple gauge group $\mathcal{G}$ and n_f chiral superfields transforming as the $\mathbf{r}$ representation of $\mathcal{G}$, expressed in terms of the gauge spinor supermultiplet W_α^A, and the chiral matter superfields Z_a^r, where A is the adjoint index of the gauge group, and r is that of the $\mathbf{r}$ representation. The index $a=1,2\ldots n_f$ spans the flavors, and α is the Weyl index of the Lorentz group.

With no superpotential, this theory is invariant under a global group $SU(n_f)\times U(1)_R$, where R is the non-anomalous R-symmetry. The gauge superfield has $R=1$ by convention, while the R value of the chiral superfields, determined by requiring the absence of the mixed $R\mathcal{G}\mathcal{G}$ anomaly, reads

$$R(Z)=1-\frac{c_A}{c_r}\frac{1}{n_f}$$

where c_A is the Dynkin index of the adjoint representation, c_r that of the matter. This theory has global anomalies. The total R anomaly (mixed gravitational anomaly) is independent of the number of flavors

$$<R>=\frac{1}{c_r}(d_A c_r - c_A d_r)\ ,$$

where $d_{A,r}$ is the dimension of the adjoint (matter) representation. In terms of $\hat{d}_r\equiv\frac{c_A}{c_r}d_r$, the other global R-anomalies are given by

$$<R^3>=d_A-\frac{\hat{d}_r^3}{(d_r n_f)^2}\ ;\qquad <RSU(n_f)^2>=-\frac{\hat{d}_r}{n_f}\ .$$

Finally the $SU(n_f)$ anomaly is just the dimension of the matter representation,

$$<SU(n_f)^3>=d_r\ .$$

They are not all independent since

$$<R^3>-<R>+n_f<RSU(n_f)^2>(1-\left(\frac{<RSU(n_f)^2>}{<SU(n_f)^3>}\right)^2)=0\ .$$

One can form an infinite number of gauge invariant holomorphic polynomials out of the W and Z fields, but they are not all independent. The independent invariants span a finite

dimensional space. In a supersymmetric theory, it is a superpace, where the bosons contain an even number of W's, and the fermions an odd-number. These invariants transform covariantly under the global symmetry $SU(n_f)$ represented by the traceless part of the operators

$$T_b{}^a = Z_b^r \frac{\partial}{\partial Z_a^r} \ ,$$

suitably summed over the gauge index to make it gauge invariant. The trace part generate a $U(1)$ symmetry. Similarly the left-handed $SU(2)_L$ Lorentz group is generated by the traceless part of

$$S_\beta{}^\alpha = W_\beta^A \frac{\partial}{\partial W_\alpha^A} \ .$$

Its trace part generates another $U(1)$. Only one linear combination of these $U(1)$s is non-anomalous. Let us introduce the gauge-invariant fermionic operators

$$\mathcal{Q}_\alpha^{a_1\cdots a_p} \equiv W_\alpha^A C_A^{r_1\cdots r_p} \frac{\partial}{\partial Z_{a_1}^{r_1}} \cdots \frac{\partial}{\partial Z_{a_p}^{r_p}} \ ,$$

where C is the appropriate Clebsch-Gordan coefficient to make an invariant, so that the adjoint is contained in the p-fold product of the matter representation. Its conjugate is given by

$$\overline{\mathcal{Q}}_{a_1\cdots a_p}^\alpha \equiv \frac{\partial}{\partial W_\alpha^A} C_A^{r_1\cdots r_p} Z_{a_1}^{r_1} \cdots Z_{a_p}^{r_p} \ .$$

Their anticommutator is generally complicated. Since the W are fermion operators, the number of such operators that appear by the use of further commutators is finite; in this sense these are like W-algebras. For any gauge group, we can build one representation of this algebra, since the polynomial

$$W^A \cdot W^A \ ,$$

which enters in the Lagrangian density always exists. It is annihilated by the lowering operator $\mathcal{Q}$, but by acting on it with the raising ladder operator $\overline{\mathcal{Q}}$, we generate the invariant fermion

$$F_{\alpha\ a_1\cdots a_p} = W_\alpha^A C_A^{r_1\cdots r_p} Z_{a_1}^{r_1} \cdots Z_{a_p}^{r_p} \ .$$

A further application generates the boson invariant

$$B_{a_1\cdots a_p b_1\cdots b_p} = C_A^{r_1\cdots r_p} C_A^{s_1\cdots s_p} Z_{a_1}^{r_1}\cdots Z_{a_p}^{r_p} Z_{b_1}^{s_1}\cdots Z_{b_p}^{s_p} \ .$$

Its structure depends on the gauge group. This algebraic structure may be enlarged considerably by gauge invariant differential operators of order $p-k$, containing k chiral matter fields.

For a special range of the number of flavors, Seiberg[1] has shown these theories to be quantum equivalent to similar theories with different gauge groups, but with the same global symmetries and anomalies. In particular, they must have the same W-superalgebras, realized in terms of the fields in the dual theory. They corresponds to inequivalent constructions of the superalgebra, which might shed some light on the nature of duality.

An infrared fixed point[2] is suggested by the renormalization group flow. The general expression for the β-function is

$$\beta = -\frac{g^3}{16\pi^2}\frac{3c_A - n_f c_r + n_f c_r \gamma(g^2)}{1 - 2c_A \frac{g^2}{16\pi^2}} \ ,$$

where γ is the anomalous dimension, given to lowest order by

$$\gamma = -\frac{g^2}{8\pi^2}\frac{d_A}{d_r} c_r + \cdots \ .$$

If the sign of the two-loop beta function is opposite that of the one-loop, there exists the possibility of an infrared fixed point at which the theory is superconformal invariant, and the anomalous dimension is given by

$$\gamma_* = 1 - 3\frac{c_A}{c_r n_f} \ .$$

Superconformal invariance requires that the dimensions are bounded by the R value of the operators $D \geq 3|R|/2$, with the bound saturated by primary operators. Let us now examine several examples, corresponding to the different values of p.

• $\mathbf{p = 1}$. In this case, the matter content is just n_f matter superfields transforming as the adjoint representation. The theory is asymptotically free as long as $n_f \leq 2$. In this simple case, the ladder operators are linear, and simply given by

$$\mathcal{Q}^a_\alpha \equiv W^A_\alpha \frac{\partial}{\partial Z^A_a} \ ; \qquad \overline{\mathcal{Q}}^\alpha_a \equiv \frac{\partial}{\partial W^A_\alpha} Z^A_a \ .$$

Their anticommutator is

$$\{\mathcal{Q}^a_\alpha, \overline{\mathcal{Q}}^\beta_b\} = \delta_\alpha{}^\beta T_b{}^a + \delta_b{}^a S_\alpha{}^\beta + \delta_\alpha{}^\beta \delta_b{}^a (\frac{1}{2} S + \frac{1}{n_f} T) \ ,$$

where

$$T_a{}^b = Z^A_a \frac{\partial}{\partial Z^A_b} - \frac{1}{n_f} \delta_a{}^b T$$

generate the Lie algebra $SU(n_f)$

$$[T_b{}^a, T_d{}^c] = \delta_b{}^c T_d{}^a - \delta_d{}^a T_b{}^c \ ,$$

with one $U(1)$ generated by

$$T = Z^A_a \frac{\partial}{\partial Z^A_a} \ .$$

Similarly, the generators of the left-handed Lorentz algebra $SU(2)_L$ are

$$S_\alpha{}^\beta = W^A_\alpha \frac{\partial}{\partial W^A_\beta} - \frac{1}{2} \delta^\beta_\alpha S \ ,$$

where S generates a $U(1)$,

$$S = W^A \frac{\partial}{\partial W^A} \ .$$

summing over the spinor indices. These are not unitary, as they act only on the analytic invariants. Note that only one combination of the two $U(1)$s appears in the anticommutator; it corresponds to that with a vanishing supertrace. These operators generates a well-known graded Lie algebra, $SU(2/n_f)$ in Kàc's classification. In general the $U(1)$ that appears in the algebra is not the same as the non-anomalous $U(1)_R$ of the gauge theory. However, for $n_f = 3$, the two coincide. At that point, the β-function vanishes, and the

theory may be understood in terms of the $N = 4$ supersymmetric gauge theory[3]. For three flavors, there can be a superpotential cubic in the three matter fields, using the structure functions of the gauge group, and the Levi-Civitá symbol for the flavor $SU(3)$.

Their simplest representation contains (WW, WZ, ZZ), which links the gauge kinetic term to the "meson" of the theory. One can generate other higher dimensional representations by acting on the order-N invariants of the algebra, $(Z)^N$. There are as many as the rank of the gauge group. It is interesting that the graded Lie algebra contains no information on the nature of the underlying gauge group. Since the duals of theories with two adjoints are not generally known, we hope that the presence of this algebra may prove relevant in their determination. In particular, this algebra must be present in any theory with adjoint matter[4], and must be built in terms of the fields in the electric and magnetic theories. In particular, the classical algebra $SU(2/1)$ will be present in all theories with $N = 2$ supersymmetry, but since its associated $U(1)$ is anomalous, it may not be directly relevant to the study of the quantum moduli space.

• $\mathbf{p = 2}$. An example is $\mathcal{G} = \mathbf{SO(n_c)}$, with $\mathbf{n_f}$ superfields in the fundamental $\mathbf{n_c}$. The adjoint is a second rank antisymmetric tensor with dimension $d_A = n_c(n_c - 1)/2$, $c_r = 1$, and $c_A = n_c - 2$. There may be[1] an infrared fixed point in the renormalization group flow if

$$\frac{3}{2} < \frac{n_f}{(n_c - 2)} < 3 \ .$$

The matter superfields Z^i_a have $R = 1 - (n_c - 2)/n_f$, the gauge superfield $W^{[ij]}_\alpha$ which contains the gauge bosons and gauginos has $R = 1$, i, j denote the $SO(n_c)$ vector indices, and a the $SU(n_f)$ index. With these, one can construct many gauge invariant chiral superfields, starting with the Lorentz singlet kinetic term

$$W^{[ij]} \cdot W^{[ij]} \ .$$

With matter superfields only, we have the flavor-symmetric "mesons",

$$I_{(ab)} = \delta_{ij} Z^i_a Z^j_b \ ,$$

with $R = 2 - 2(n_c - 2)/n_f$, and a twice antisymmetric spinor invariant

$$F_{\alpha[ab]} = W^{[ij]}_\alpha Z^i_a Z^j_b \ ,$$

with $R = 2(2n_f - n_c + 2)/n_f$.

There is another class of "topological" invariant composites constructed with the flavor Levi-Cività symbol: two bosons, one n_c-antisymmetrized composite

$$I_{[a_1 \cdots a_{n_c}]} = \epsilon^{i_1 i_2 \cdots i_{n_c}} Z^{i_1}_{a_1} Z^{i_2}_{a_2} \cdots Z^{i_{n_c}}_{a_{n_c}} \ ,$$

with $R = n_c(1 - (n_c - 2)/n_f)$, and one $(n_c - 4)$-antisymmetrized composite

$$I_{[a_1 \cdots a_{n_c - 4}]} = \epsilon^{i_1 i_2 \cdots i_{n_c}} W^{[i_1 i_2]} \cdot W^{[i_3 i_4]} Z^{i_5}_{a_1} Z^{i_6}_{a_2} \cdots Z^{i_{n_c}}_{a_{n_c - 4}} \ ,$$

with $R = 2 + (n_c - 4)(n_f - n_c + 2)/n_f$. There is also one topological spinor invariant, the $(n_c - 2)$-antisymmetric flavor tensor

$$F_{\alpha[a_1 a_2 \cdots a_{n_c - 2}]} = \epsilon^{i_1 i_2 \cdots i_{n_c}} W^{[i_1 i_2]}_\alpha Z^{i_3}_{a_1} Z^{i_4}_{a_2} \cdots Z^{i_{n_c}}_{a_{n_c - 2}} \ ,$$

with $R = 1 + (n_c - 2)(n_f - n_c + 2)/n_f$. They are primary invariants[1]. When normalized to the meson invariant, they span a finite-dimensional compact superspace, the *superorbit* space of the theory.

To build the graded W-algebra associated with this theory, we start with the lowering ladder operators

$$\mathcal{Q}^{ab}_\alpha = W^{ij}_\alpha \frac{\partial}{\partial Z^i_a} \frac{\partial}{\partial Z^j_b} \ ,$$

and their conjugates

$$\overline{\mathcal{Q}}^\alpha_{ab} = Z^i_a Z^j_b \frac{\partial}{\partial W^{ij}_\alpha} \ ,$$

which act as raising operators. These transform as the twice-antisymmetrized repesentation of the flavor $SU(n_f)$, and its conjugate, with $R = \pm(n_f - n_c + 2)/n_f$. Both annihilate the quadratic meson invariant I_{ab}, which forms a trivial representation of the algebra.

By acting the raising operator on the pure gauge invariant WW, we generate the fermion invariant $F_{\alpha\ [ab]}$. Another application yields the *square* of the meson invariant. This is the representation of invariants that always exists. Since it contains the square of an invariant, it cannot be viewed as fundamental. It also shows that these ladder operators do not commute with the Lagrangian.

The primary (topological) invariants form a different representation of the superalgebra. We find symbolically that

$$\mathcal{Q} I_{[a_1 \cdots a_{n_c}]} = F_{[a_1 \cdots a_{n_c-2}]} \ ; \qquad \mathcal{Q} F_{[a_1 \cdots a_{n_c-2}]} = I_{[a_1 \cdots a_{n_c-4}]} \ ; \qquad \mathcal{Q} I_{[a_1 \cdots a_{n_c-4}]} = 0 \ ,$$

the last obtained by symmetry properties. The Lie algebra generators of the global $SU(n_f) \times SU(2)_L \times U(1) \times U(1)$ are generated by

$$T_a{}^b = Z_a^j \frac{\partial}{\partial Z_b^j} \ ; \qquad S_\alpha{}^\beta = W_\alpha^{ij} \frac{\partial}{\partial W_\beta^{ij}} \ .$$

These are not unitary, as they act only on the analytic invariants. Under these, the fermionic ladder operators transform as expected,

$$[T_c{}^d, \mathcal{Q}_\alpha^{ab}] = \delta^d{}_a \mathcal{Q}_\alpha^{cb} - \delta^d{}_b \mathcal{Q}_\alpha^{ac} \ ,$$

$$[S_\alpha{}^\beta, \mathcal{Q}_\gamma^{ab}] = \delta^\beta{}_\gamma \mathcal{Q}_\alpha^{ab} \ .$$

The anticommutator of the raising and lowering operators is more complicated; it contains not only a part linear and quadratic in the Lie generators, but also new operators

$$\begin{aligned}[\mathcal{Q}_\alpha^{ab}, \overline{\mathcal{Q}}_{cd}^\beta] = \delta_{cd}^{ab} S_\alpha{}^\beta - \delta_\alpha^\beta (T_c{}^a T_d{}^b - T_d{}^a T_c{}^b + \delta^a{}_c T_d{}^b - \delta^a{}_d T_c{}^b) + \\ + \delta_c{}^a U_{\alpha d}^{\beta b} - \delta_d{}^a U_{\alpha c}^{\beta b} + \delta_d{}^b U_{\alpha c}^{\beta a} - \delta_c{}^b U_{\alpha d}^{\beta a} \ ,\end{aligned}$$

where

$$\delta^{ab}_{cd} \equiv \delta^a{}_c \delta^b{}_d - \delta^b{}_c \delta^a{}_d \ ,$$

and

$$U^{\beta a}_{\alpha c} = W^{im}_{\alpha} \frac{\partial}{\partial W^{mj}_{\beta}} Z^j_b \frac{\partial}{\partial Z^i_a} \ ,$$

are the new operators. Their commutators in turn yield higher order operators and so on, up to a power related to the number of distinct fermions, in this case, $2n_c(n_c - 1)$.

We have not separated the traces of $T_b{}^a$ and $S^{\beta}{}_{\alpha}$, which correspond to two $U(1)$ symmetries. They are present in the underlying gauge theory, but one is anomalous. We have not been able to determine algebraically the condition that singles out the relevant linear combination of these two $U(1)$s.

While complicated, this algebra may shed some light on the origin of the Seiberg duals. The dual of this theory is of the same type with a $SO(\tilde{n}_c)$ gauge group, represented by the gauge superfield $\tilde{W}^{\tilde{i}\tilde{j}}_{\alpha}$, and n_f matter fields $\tilde{Z}^{\tilde{i}\ a}$, transforming contragrediently under the flavor goup. Thus the same algebra can be constructed out of the tilded fields, with the important difference that the flavor properties of the raising and lowering operators are interchanged.

The same primary invariants are built out of the tilded fields: a $\tilde{n}_c$-antisymmetric flavor tensor

$$\tilde{I}^{[a_1 \cdots a_{\tilde{n}_c}]} = \epsilon^{\tilde{i}_1 \tilde{i}_2 \cdots \tilde{i}_{\tilde{n}_c}} \tilde{Z}^{a_1 \tilde{i}_1} \tilde{Z}^{a_2 \tilde{i}_2} \cdots \tilde{Z}^{a_{\tilde{n}_c} \tilde{i}_{\tilde{n}_c}} \ .$$

Anomaly cancellation requires that

$$\frac{\tilde{n}_c - 2}{n_f} + \frac{n_c - 2}{n_f} = 1 \ ,$$

so that this invariant has exactly the same quantum numbers as $I_{[a_1 \cdots a_{n_c - 4}]}$, the invariant that contains two gauge supermultiplets in the original theory. This suggests that duality

corresponds to a conjugation of this algebra, interchanging the role of raising and lowering operators.

• $\mathbf{p = 3}$. An example is $\mathcal{G} = \mathbf{E_6}$, with $\mathbf{n_f}$ chiral superfields transforming as the $\mathbf{27}$. Since $c_A = 24$ and $c_r = 6$, and $\hat{d} = 4$, the matter superfields have $R_m = 1 - 4/n_f$. The theory is asymptotically free as long as $n_f < 12$, and since the two-loop coefficient is proportional to $108 - 22n_f$, it may have an infrared fixed point when $n_f \geq 5$. The values of its anomalies are

$$< R >= \ -30 \ ; \qquad < R^3 >= \ 78 - \frac{(12)^3}{n_f^2} \ ;$$
$$< RSU(n_f)^2 >= \ -\frac{108}{n_f} \ ; \qquad < SU(n_f)^3 >= \ 27 \ .$$

The lowest invariant that can be built out of the $(\mathbf{27}, \mathbf{n}_f)$ is the symmetric cubic invariant, $\mathbf{S^{(rst)}}$. We expect it to be present in the dual theory either as a fundamental field or as a construct of the fields in the dual theory. Its R value is $3(1 - 4/n_f)$. Assume that $\mathbf{S^{(rst)}}$ is a fundamental field of the dual theory; its contributions to the anomalies are[5]

$$\begin{aligned} < R >&= \frac{1}{3}(n_f + 1)(n_f + 2)(n_f - 6) \ , \\ < R^3 >&= \frac{4}{3n_f^2}(n_f + 1)(n_f + 2)(n_f - 6)^3 \ , \\ < RSU(n_f)^2 >&= \frac{(n_f + 2)(n_f + 3)(n_f - 6)}{n_f} \ , \\ < SU(n_f)^3 >&= \frac{(n_f + 3)(n_f + 6)}{2} \ . \end{aligned}$$

The only obvious match is when $n_f = 6$: the anomaly of $\mathbf{S^{(rst)}}$ is equal 54, with $R = 3(1 - 4/6) = 1$, and its fermions do not contribute to three of the global anomalies. On the other hand, the excess $SU(6)^3$ anomaly is 27, suggesting that the dual theory is also an E_6 gauge theory with its $\mathbf{27}$ transforming as the $\overline{\mathbf{n}}_\mathbf{f}$, with the matter content

$$\tilde{\mathbf{Z}}_r \sim (\mathbf{27}, \overline{\mathbf{6}}) \ \text{ and } \mathbf{S^{(rst)}} \sim (\mathbf{1}, \mathbf{35}) \ ,$$

and superpotential

$$W = \tilde{\mathbf{Z}}_r \tilde{\mathbf{Z}}_s \tilde{\mathbf{Z}}_t \mathbf{S}^{(\mathbf{rst})} \ .$$

This is exactly the same structure as the self dual point of $SO(n_c)$ with $n_f = 2(n_c - 2)$. Besides the anomaly matching conditions, there is corroborating evidence from the "baryons" of the theory: there is a sixth order composite invariant with mixed flavor symmetry[6]

$$(\mathbf{27}, \square)^6 \sim (\mathbf{1}, \begin{smallmatrix}\square&\square\\\square&\square\end{smallmatrix}) \ .$$

For duality to hold, it must be manufactured equally well by the electric as by the magnetic fundamental fields.

The fields of the magnetic theory transform as the conjugate flavor representation, the same sixth order invariant of the magnetic theory will give six contragredient flavor indices, which can be transformed to six "lower" flavor indices by means of two Levi-Cività symbols.

For a larger value of n_f, say $n_f = 7$, the same procedure yields a flavor invariant that transforms as an eighth order invariant in flavor space. Thus the dual theory for E_6 with seven flavors must contain the flavor combination given by the Young tableau

$$\begin{smallmatrix}\square&\square\\\square&\square\\\square&\square\\\square&\square\end{smallmatrix} \ .$$

We have not succeeded in finding this dual.

The gauge invariant fermionic ladder operator is a third order operator, since in E_6, the adjoint representation appears only in the cubic product of **27**, with mixed flavor symmetry[7]

$$\frac{\partial}{\partial W^{ijk}} Z^i_a Z^j_b Z^k_c \sim (\mathbf{1}, \begin{smallmatrix}\square&\square\\\square&\end{smallmatrix}) \ ,$$

where i, j, k are **27** indices. We can apply this operator to the gauge condensate and obtain

a fermionic chiral superfields

$$W^{ijk} Z^i_a Z^j_b Z^k_c \sim (\mathbf{1}, \begin{smallmatrix}\square\\ \square\end{smallmatrix}) \ .$$

Another application yields sixth-order invariants. It may be the square of the "meson" field, as in the orthogonal group case, but it most likely contains the independent sixth-order "baryon" invariant.

We can consider more general cases, such as $p = 4$, an example of which is an $SO(10)$ theory with matter in its spinor representation.

I wish to thank the Aspen Center for Physics where this note was put together, and for the many useful remarks of its participants in particular P. Binétruy, E. Dudas, G. Moore, J. Patera, C. Savoy, and B. Zumino.

References

1) For a review see K. Intriligator and N. Seiberg, *Lectures on Supersymmetric Gauge Theories and Electric-Magnetic Duality*, hep-th/9509066, and references herein. N. Seiberg, *Nucl. Phys.* **B447** (1995) 95.

2) T. Banks and A. Zaks, *Nucl. Phys.* **B196** (1982) 189. M.A. Shifman and A.I. Vainstein, *Nucl. Phys.* **B359** (1991) 571.

3) I thank M. Strassler for this remark.

4) K. Intriligator, R.G. Leigh, M.J. Strassler, *Nucl. Phys.* **B456** (1995) 567.

5) J. Patera, and R.T. Sharp, *J. Math. Phys.* **22** (1981) 2352.

6) S. Giddings, private communication. The search for the dual of E_6 was started in a collaboration with A. Nelson and S. Giddings.

7) J. Patera, private communication.